arXiv:1610.09016v1 [math.HO] 25 Oct 2016

On the weighted Gauß–Radau Quadrature

Sascha Trostorff & Marcus Waurick*

October 31, 2016

Abstract

In this short note, we collect some facts on the weighted Gauß-Radau quadrature. In particular, we focus on the location of the Gauß–Radau points being a continuous function of the L^1-weighting function.

AMS subject classification (2010): 65D32, 65D30, 65D05
Key words: numerical quadrature, weighted integrals, Gauß–Radau points

1 On the Gauß–Radau Quadrature

In this technical paper we shall gather some results on the right-sided Gauß–Radau quadrature, which are well-known in principle, but are collected for the convenience of the reader, in particular we refer to [1]. For this, we introduce a set of weighting functions:

$$W := \{w \in L^1(-1,1) : w > 0 \text{ a.e.}\}.$$

Note that the bilinear form

$$\langle \cdot, \cdot \rangle_w \colon D \times D \ni (f,g) \mapsto \int_{-1}^{1} f(x)g(x)w(x)\,\mathrm{d}x$$

introduces a scalar product on its natural domain

$$D := \{f \in L^1_{\mathrm{loc}}(-1,1); \int_{(-1,1)} |f(x)|^2 w(x)\,\mathrm{d}x < \infty\}.$$

Throughout, let $q \in \mathbb{N}$.

*Institut für Analysis, TU Dresden, Dresden, Germany, e-mail: sascha.trostorff@tu-dresden.de & Department of Mathematical Sciences, University of Bath, Bath, UK. e-mail: m.waurick@bath.ac.uk

Definition. Let $w \in W$. A pair $(\omega, r) = ((\omega_j)_{j\in\{0,\dots,q\}}, (r_j)_{j\in\{0,\dots,q\}}) \in \mathbb{R}^{q+1} \times [-1,1]^{q+1}$ is called *(right-sided) w-Gauß–Radau quadrature (of order q)*, if $-1 \le r_0 \le r_1 \le \cdots \le r_q = 1$ and for all $p \in \mathcal{P}_{2q}(-1,1)$ we have

$$\int_{-1}^{1} p(x)w(x)\,\mathrm{d}x = \sum_{j=0}^{q} \omega_j p(r_j).$$

Theorem 1.1. *Let $w \in W$. Then there exists a unique (right-sided) w-Gauß–Radau quadrature (ω, r) of order q. The weights ω satisfy $0 < \omega_j \le \int_{(-1,1)} w(x)\,\mathrm{d}x$ for $j \in \{0,\dots,q\}$ and $r_0 > -1$.*

The result being well-known we only refer to [2] for the corresponding result for Gauß-quadrature rule. Note that the methods in [2] can straightforwardly be adopted to apply to the Gauß–Radau quadrature, if one considers the inner product $\langle\cdot,\cdot\rangle_{\tilde{w}}$ instead of $\langle\cdot,\cdot\rangle_w$, where for $w \in W$ we put $\tilde{w}\colon x \mapsto (1-x)w(x)$ and observe $\tilde{w} \in W$.

Next, we address the continuous dependence of the Gauß–Radau points on the weighting function.

Theorem 1.2. *The mapping*

$$(W, \|\cdot\|_{L^1(-1,1)}) \to \mathbb{R}^{q+1} \times (-1,1]^{q+1}, w \mapsto (\omega(w), r(w)) \tag{1}$$

is continuous, where $(\omega(w), r(w))$ denotes the w-Gauß–Radau quadrature.

Proof. Let $(w_n)_{n\in\mathbb{N}}$ be a sequence in W and $w \in W$ such that $w_n \to w$ in $L^1(-1,1)$. By definition and by Theorem 1.1,

$$(\omega(w_n), r(w_n)) \in [0, \sup_{k\in\mathbb{N}} \|w_k\|_{L^1(-1,1)}]^{q+1} \times [-1,1]^{q+1}$$

for all $n \in \mathbb{N}$. Thus, there exists a convergent subsequence for which we reuse the name with limit $(\overline{\omega}, \overline{r})$. Note that $\overline{r}_q = 1$. Next, let $p \in \mathcal{P}_{2q}(-1,1)$. Then, for $n \in \mathbb{N}$, we obtain

$$\begin{aligned}\int_{-1}^{1} pw &= \lim_{n\to\infty} \int_{-1}^{1} pw_n \\ &= \lim_{n\to\infty} \sum_{j=0}^{q} \omega(w_n)_j p(r(w_n)_j) \\ &= \sum_{j=0}^{q} \overline{\omega}_j p(\overline{r}_j).\end{aligned}$$

Hence, by Theorem 1.1, we infer $(\overline{\omega}, \overline{r}) = (\omega(w), r(w))$, which eventually implies the assertion. □

Corollary 1.3. *For $\tau \in \mathbb{R}$ denote $w_\tau \colon x \mapsto \exp(-\rho\tau(x+1))(\in W)$ and let $(\omega^{(\tau)}, r^{(\tau)})$ be the w_τ-Gauß–Radau quadrature. For $\tau \in \mathbb{R}$, let $\chi_\tau \in \mathcal{P}_{q+1}(-1,1)$ such that*

$$\chi_\tau(r_j^{(\tau)}) = 0 \quad (j \in \{0,\dots,q\}),\ \chi_\tau(-1) = 1.$$

Then, for every compact set $K \subset \mathbb{R}$, we have

$$\sup_{\tau\in K} \int_{-1}^{1} \chi_\tau^2 w_\tau < \infty.$$

Proof. Assume by contradiction that there exists $(\tau_n)_n$ convergent to some τ with the property

$$\int_{-1}^{1} \chi_{\tau_n}^2 w_{\tau_n} \to \infty.$$

Using Theorem 1.2, we compute for $n \in \mathbb{N}$

$$\begin{aligned}
\int_{-1}^{1} \chi_{\tau_n}^2 w_{\tau_n} &= \int_{-1}^{1} \prod_{j=0}^{q} \Big(\frac{x - r_j^{(\tau_n)}}{-1 - r_j^{(\tau_n)}}\Big)^2 w_{\tau_n}(x)\,\mathrm{d}x \\
&\le \int_{-1}^{1} \prod_{j=0}^{q-1} \Big(\frac{x - r_j^{(\tau_n)}}{-1 - r_j^{(\tau_n)}}\Big)^2 w_{\tau_n}(x)\,\mathrm{d}x \\
&= \sum_{\ell=0}^{q} \omega_\ell^{(\tau_n)} \prod_{j=0}^{q-1} \Big(\frac{r_\ell^{(\tau_n)} - r_j^{(\tau_n)}}{1 + r_j^{(\tau_n)}}\Big)^2 \\
&= \omega_q^{(\tau_n)} \prod_{j=0}^{q-1} \Big(\frac{1 - r_j^{(\tau_n)}}{1 + r_j^{(\tau_n)}}\Big)^2 \\
&\to \omega_q^{(\tau)} \prod_{j=0}^{q-1} \Big(\frac{1 - r_j^{(\tau)}}{1 + r_j^{(\tau)}}\Big)^2 \quad (n \to \infty).
\end{aligned}$$

But,

$$0 \le \omega_q^{(\tau)} \prod_{j=0}^{q-1} \Big(\frac{1 - r_j^{(\tau)}}{1 + r_j^{(\tau)}}\Big)^2 = \int_{-1}^{1} \prod_{j=0}^{q-1} \Big(\frac{x - r_j^{(\tau)}}{-1 - r_j^{(\tau)}}\Big)^2 w_\tau(x)\,\mathrm{d}x < \infty,$$

which contradicts the assumption. □

Let us introduce for a bounded interval $I \subseteq \mathbb{R}$ the mapping

$$\begin{aligned}
\phi_I \colon (-1,1) &\to I, \\
x &\mapsto \frac{a+b}{2} + \frac{b-a}{2} x,
\end{aligned}$$

where $a \coloneqq \inf I$, $b \coloneqq \sup I$. Further, we set $|I| \coloneqq b - a$.

Corollary 1.4. *For $\tau \in \mathbb{R}$ let χ_τ be as in Corollary 1.3. Let $K \geq 0$. Then*

$$\sup_{I \subseteq \mathbb{R} \text{ interval}, |I| \leq K} \frac{1}{|I|} \int_I \big(\chi_{|I|}(\phi_I^{-1}(t))\big)^2 e^{-2\rho(t-\inf I)} \,\mathrm{d}t < \infty.$$

Proof. For $I = (a, b) \subseteq \mathbb{R}$ we compute

$$\begin{aligned}
&\frac{1}{b-a} \int_a^b \big(\chi_{|I|}(\phi_I^{-1}(t))\big)^2 e^{-2\rho(t-a)} \,\mathrm{d}t \\
&= \frac{1}{b-a} \int_{-1}^1 \big(\chi_{|I|}(x)\big)^2 e^{-2\rho(\phi_I(t)-a)} \phi_I'(x) \,\mathrm{d}x \\
&= \frac{1}{2} \int_{-1}^1 \big(\chi_{|I|}(x)\big)^2 e^{-\rho((b-a)(x+1))} \,\mathrm{d}x.
\end{aligned}$$

Hence, the assertion follows from Corollary 1.3. □

The next corollary is concerned with the lowest Gauß–Radau point for different weights:

Corollary 1.5. *For $\tau \in \mathbb{R}$ let w_τ and $(\omega^{(\tau)}, r^{(\tau)})$ be given as in Corollary 1.3. Let $T > 0$. Then there exists $c > 0$ such that for all intervals $I \subseteq \mathbb{R}$ with $|I| \leq T$ and $0 \leq \tau \leq T$ we have*

$$\phi_I(r_0^{(\tau)}) - \inf I \geq c|I|.$$

Proof. We observe that

$$\mathbb{R} \ni \tau \mapsto \big(t \mapsto e^{-\rho\tau(t+1)}\big) \in (W, \|\cdot\|_{L^1(-1,1)})$$

is continuous. Hence, the set

$$\{\big(t \mapsto e^{-\rho\tau(t+1)}\big); \tau \in [0, T]\} \subseteq (W, \|\cdot\|_{L^1(-1,1)})$$

is compact. Thus, by the continuous dependence of the Gauß–Radau point on the weighting function (see (1)), we obtain that

$$\{(\omega^{(\tau)}, r^{(\tau)}); \tau \in [0, T]\} \subseteq \mathbb{R}^{q+1} \times (-1, 1]^{q+1}$$

is compact, as well. In particular, there exists $c > 0$ with the property $r_0^{(\tau)} - (-1) \geq c$. Hence, we obtain for all $\tau \in [0, T]$ and intervals $I \subseteq \mathbb{R}$ with $|I| \leq T$

$$\phi_I(r_0^{(\tau)}) - \inf I = \phi_I(r_0^{(\tau)}) - \phi_I(-1) = |I|(r_0^{(\tau)} - (-1)) \geq c|I|. \qquad \square$$

References

[1] S. Franz, S. Trostorff, M. Waurick. Numerical Methods for Changing Type Systems. Preprint, TU Dresden, 2016

[2] G. Lube Problemstellung. Orthogonale Polynome Online-manuscript: `https://lp.uni-goettingen.de/get/text/1275` and `https://lp.uni-goettingen.de/get/text/1276` available at 25/10/2016, Georg-August-University Göttingen, 2004.